



Taxonomic Paper

# The Coreidae of Honduras (Hemiptera: Coreidae)

Carlos A Linares<sup>‡</sup>, Jesus Orozco<sup>‡</sup>

<sup>‡</sup> Insect Collection, Zamorano University, Zamorano, Honduras

Corresponding author: Jesus Orozco ([jorozco@zamorano.edu](mailto:jorozco@zamorano.edu))

Academic editor: Laurence Livermore

Received: 04 Apr 2017 | Accepted: 02 Jun 2017 | Published: 05 Jun 2017

Citation: Linares C, Orozco J (2017) The Coreidae of Honduras (Hemiptera: Coreidae). Biodiversity Data Journal 5: e13067. <https://doi.org/10.3897/BDJ.5.e13067>

## Abstract

### Background

Coreidae bugs are mostly sap-sucking insects feeding on a variety of plants. Despite their abundance and economic importance in Honduras there is little information on the species, their distribution and affected crops. Since knowledge of pest species allows for better management of crops, we aimed to document the diversity of this economically important group. Specimens from four entomological collections in Honduras were studied and an exhaustive search of all available literature was conducted.

### New information

A total of 2,036 insects were examined. The fauna of Honduran coreids is now composed of 68 species. Nineteen species are recorded for the country for the first time and 17 species were found only in literature. Little is known about the biology and economic importance of most of the species.

## Keywords

Taxonomy, diversity, agriculture, pest, Central America.

## Introduction

Bugs of the Coreidae family are primarily phytophagous insects that feed on plants sucking sap from branches, leaves, flowers and fruits. Many coreids are known pests of ornamentals and crops that can, at times, cause serious damage or even total loss (Henry 2009). Due to their feeding habits, they can cause malformations, rotting, discoloration, and also abortion of the fruit and poor formation of the seed (Mitchell 2000).

Honduras' economy is largely depending on agriculture. Given that pest control depends heavily on proper species identification and management, knowledge of the species becomes paramount for the economy. Despite this, little information on insects from the country is available.

Evans and Halbert (2007) found 26 new aphids in Honduras out of the 46 species known for the country. In Coleoptera, Turnbow et al. (2003b), found that of the 626 species of Honduran cerambycids, 364 were new country records. Similarly, Turnbow et al. (2003a), found 78 bruchids new to the country of the 148 known species. We estimate that at least half of the species of insects in Honduras are known unknowns: species already described that are not recorded for the country. This, together with the relative lack of in-country taxonomists poses a difficulty for, among other things, crop management. It is impossible to know the pest status of an unknown insect.

This work is an effort to provide comprehensive information on the coreid species from Honduras, including their geographic distribution by department and the plants they are known to feed on.

## Materials and methods

Specimens from the following entomological collections in Honduras were examined:

- Colección Entomológica del Centro Universitario Regional del Litoral Atlántico, Ceiba (CURLA).
- Colección Entomológica de la Escuela Nacional de Ciencias Forestales, Siguatepeque (CEEFF).
- Museo Entomológico de la Universidad Nacional Autónoma de Honduras, Tegucigalpa (UNAH).
- Zamorano Insect Collection, Zamorano University, Zamorano (EAPZ).

The material was identified by comparison with a reference collection at EAPZ and by using available keys. Label information containing host, date of collection, and distribution was recorded. Additional species information was gathered from the literature.

Geographic and temporal distribution of adults in Honduras as well as biological information for all species was obtained from label data. For the new country records the known distribution, outside Honduras, according to the available literature is included.



## List of Honduran Coreids

### *Acanthocephala alata* (Burmeister, 1835)

**Distribution:** Atlántida, Comayagua, Francisco Morazán, and Yoro.

**Notes:** Specimens examined: 115 (CEEF, CURLA, EAPZ, UNAH).

Temporal distribution: May–September.

Hosts: *Phaseolus vulgaris* L. (beans) (EAPZ); *Annona muricata* L. (soursop) (Hernández and Pinzón 2015), and *Jatropha curcas* L. (Barbados nut) (Grimm and Maes 1997).

Attacks the fruits of *Annona muricata* L. causing their fall (Hernández and Pinzón 2015).

### *Acanthocephala declivis* (Say, 1832)

**Distribution:** Atlántida, Comayagua, and Francisco Morazán.

**Notes:** Specimens examined: 13 (CEEF, CURLA, EAPZ, UNAH).

Temporal distribution: May–July.

Hosts: *Baccharis halimifolia* L. (groundsel bush), *Baccharis neglecta* Britton, and *Persea borbonia* (L.) Spreng. (redbay) (McPherson et al. 2011).

### *Acanthocephala femorata* (F., 1775)

**Distribution:** Atlántida, Comayagua, El Paraíso, Francisco Morazán, and Valle.

**Notes:** Specimens examined: 108 (CEEF, CURLA, EAPZ, UNAH).

Temporal distribution: January (Passoa 1983), May–July, October.

Hosts: *Ipomoea batatas* (L.) Lam. (sweet potato), *Citrus sinensis* (L.) Osbeck (orange), *Cucurbita pepo* L. (pumpkin) (EAPZ); *Solanum tuberosum* L. (potato) (Passoa 1983); *Helianthus annuus* L. (sunflower), *Cirsium texanum* Buckley, *Ambrosia trifida* L. (wild hemp), *Baccharis neglecta* Britton, *Chenopodium album* L. (pigweed), *Cirsium horridulum* Michx. (yellow thistle), *Ratibida columnifera* (Nutt.) Wooton & Standl. (prairie coneflower), *Sorghum halepense* (L.) Pers. (Johnson grass), *Erigeron quercifolius* Lam., *Abelmoschus esculentus* (L.) Moench (gumbo), *Parthenium* sp., *Gossypium hirsutum* L. (cotton), *Prunus persica* (L.) Batsch (peach), *Sorghastrum nutans* (L.) Nash (McPherson et al. 2011), and *Jatropha curcas* L. (Barbados nut) (Alonso and Lezcano 2014).

*Acanthocephala femorata* is parasitized by *Trichopoda pennipes* (F.), (Diptera: Tachinidae). McPherson et al. (2011) recorded *Arilus cristatus* L. (Hemiptera: Reduviidae), *Gryon floridanum* Ashmead (Hymenoptera: Scelionidae), and *Bicyrtes quadrifasciata* Say (Hymenoptera: Crabronidae) as natural enemies.

### ***Anasa bellator* (F., 1787)**

**Distribution:** Unknown in Honduras (Maes and Goellner-Scheiding 1993).

**Notes:** Temporal distribution: July–October (Maes and Goellner-Scheiding 1993).

Hosts: *Tournefortia* sp., *Zea mays* L. (corn), *Coffea arabica* L. (coffee) (Maes and Goellner-Scheiding 1993), and *Myristica fragrans* Houtt. (nutmeg) (Brailovsky 1985).

### ***Anasa capaneodes* Stål, 1862**

**Distribution:** Francisco Morazán.

**Notes:** Specimens examined: 5 (EAPZ).

Temporal distribution: May–July.

Hosts: *Tillandsia bourgaei* Baker, *Tillandsia intumescens* L. B. Sm. and *Tillandsia prodigiosa* (Lem.) Baker (Maes and Goellner-Scheiding 1993).

### ***Anasa linnavuorii* Brailovsky, 2016**

**Distribution:** Yoro (Brailovsky 2016)

**Notes:** Temporal distribution: June (Brailovsky 2016).

Hosts: Unknown (Brailovsky 2016).

### ***Anasa maculipes* Stål, 1862**

**Distribution:** Olancho (Brailovsky 1985).

**Notes:** Temporal distribution: Unknown.

Hosts: *Cucurbita pepo* L. (pumpkin), *Quercus* sp., *Buddleja americana* L., and *Ageratina adenophora* (Spreng.) R.M. King & H. Rob. (crofton-weed) (Brailovsky 1985).

### ***Anasa scorbutica* (F., 1775)**

**Distribution:** Atlántida, Choluteca, Comayagua, El Paraíso, Francisco Morazán, La Paz, Olancho, and Yoro.



**Notes:** Specimens examined: 35 (CEEF, CURLA, EAPZ, UNAH).

Temporal distribution: May–December.

Hosts: *Cucurbita argyrosperma* K. Koch, *Zea mays* L. (corn), *Cucurbita pepo* L. (pumpkin) (Passoa 1983); *Solanum lycopersicum* L. (tomato), *Lagenaria siceraria* (Molina) Standl. (bottle gourd) (Brailovsky 1985), and *Jatropha curcas* L. (Barbados nut) (Alonso and Lezcano 2014).

Brailovsky (1985) recorded *Ooencyrtus submetallicus* Howard (Hymenoptera: Encyrtidae), *Anastatus diversus* Gahan (Hymenoptera: Eupelmidae), and *Gryon carinatifrons* Say (Hymenoptera: Crabronidae) as natural enemies.

### ***Anasa trilineata* Stål, 1870**

**Distribution:** El Paraíso and Francisco Morazán.

**Notes:** NEW COUNTRY RECORD

Specimens examined: 11 (EAPZ).

Temporal distribution: May–July.

Hosts: *Sechium edule* (Jacq.) Sw. and *Zea mays* L. (corn) (EAPZ).

Known distribution: Brazil, Colombia, Costa Rica, Ecuador, Nicaragua, Paraguay, Peru, and Venezuela (Packauskas 2010).

### ***Anasa tristis* (De Geer, 1773)**

**Distribution:** Francisco Morazán.

**Notes:** Specimens examined: 6 (CEEF, EAPZ, UNAH).

Temporal distribution: May–July.

Hosts: *Cucurbita argyrosperma* K. Koch, *Cucurbita pepo* L. (pumpkin), *Citrullus vulgaris* Schrad. (watermelon), and *Cucumis melo* L. (sweet melon) (Alston and Barnhill 2008).

In the United States this species is considered as one of the important pests of pumpkin and squash. It causes necrosis on the leaves, scars on the fruits and a rapid wilting of the plant. It is parasitized by *Trichopoda pennipes* (F.) (Diptera: Tachinidae) (Alston and Barnhill 2008).

### ***Anasa uhleri* Stål, 1868**

**Distribution:** Cortés (Brailovsky 1985).

**Notes:** Temporal distribution: Unknown.

Hosts: *Cucurbita pepo* L. (pumpkin), *Opuntia streptacantha* Lem. and *Ageratina adenophora* (Spreng.) R.M. King & H. Rob. (Brailovsky 1985).

### ***Anisoscelis affinis* Westwood, 1840**

**Distribution:** Atlántida, Francisco Morazán, and Olancho.

**Notes:** Specimens examined: 40 (CEEF, CURLA, EAPZ, UNAH).

Temporal distribution: May–November.

Hosts: *Passiflora edulis* Sims (passion fruit) (EAPZ), *Passiflora quadrangularis* L. (badea) (Lerma et al. 1986), and *Solanum betaceum* Cavanilles (tree tomato) (Lucas et al. 2010).

It is considered as one of the main pests of tree tomato in Ecuador (Lucas et al. 2010).

### ***Camptischium clavipes* (F., 1803)**

**Distribution:** Atlántida.

**Notes: NEW COUNTRY RECORD**

Specimens examined: 8 (EAPZ).

Temporal distribution: February.

Hosts: *Solanum melongena* L. (eggplant) (King and Saunders 1984).

Known distribution: Argentina, Bolivia, Brazil, Guyana, Lesser Antilles, Panama, and Uruguay (Packauskas 2010).

This species sucks sap from tender shoots and fruits on eggplant plantations. It can cause decay and deformation to fruits (King and Saunders 1984).

### ***Catorhintha apicalis* (Dallas, 1852)**

**Distribution:** Distribution in Honduras unknown (Packauskas 2010).

**Notes:** Temporal distribution: June–September (Báez and Cervantes 2014).

Hosts: *Mirabilis jalapa* L., *Nolina parviflora* Kunth (Hemsl.), *Sphaeralcea* sp., and *Gossypium* sp. (Báez and Cervantes 2014).



***Catorhintha guttula* (F., 1794)**

**Distribution:** Atlántida, Choluteca, Comayagua, and La Paz.

**Notes:** Specimens examined: 1 (CURLA).

Temporal distribution: July.

Hosts: *Phaseolus vulgaris* L. (beans), *Zea mays* L. (corn) (Passoa 1983); *Mirabilis jalapa* L. (Cervantes et al. 2014); *Cirsium* sp., *Artemisia vulgaris* L., *Cucurbita pepo* L. (pumpkin), *Lyonia* sp., *Crotalaria* sp., *Glycine max* (L.) Merr. (soy), *Gossypium hirsutum* L. (cotton), *Sida* sp., *Mimosa pudica* L. *Mirabilis* sp. *Ricinus* sp., *Boerhavia* sp., *Oryza sativa* L. (rice), *Solanum lycopersicum* L. (tomato), *Theobroma* sp., and *Waltheria americana* L. (Maes and Goellner-Scheiding 1993).

***Catorhintha selector* Stål, 1860**

**Distribution:** Distribution in Honduras unknown (Brailovsky and Garcia 1987).

**Notes:** Hosts: *Gossypium herbaceum* L. (cotton), *Boerhaavia diffusa* L., *Coffea arabica* L. (coffee), *Waltheria* sp. (Maes and Goellner-Scheiding 1993), and *Mirabilis jalapa* L. (Cervantes et al. 2014).

This species is frequently associated with Nyctaginaceae (Cervantes et al. 2014).

***Cebrenis danieli* Brailovsky, 1995**

**Distribution:** Comayagua (Brailovsky 1995).

**Notes:** Temporal distribution: Unknown.

Hosts: *Neurolaena lobata* (L.) R. Br. ex Cass., *Mikania scandens* (L.) Willd., and *Verbesina* sp. (Brailovsky 1995).

***Cebreniscella exitiosa* (Brailovsky, 1984)**

**Distribution:** Distribution in Honduras unknown (Packauskas 2010).

**Notes:** Temporal distribution: Unknown.

Hosts: Unknown.

***Cebrenistella caltumae* Brailovsky, 2013**

**Distribution:** Distribution in Honduras unknown (Brailovsky 2013).

**Notes:** Temporal distribution: Unknown.

Hosts: Unknown.

***Chariesterus moestus* Burmeister, 1835**

**Distribution:** Comayagua and Francisco Morazán.

**Notes:** Specimens examined: 11 (EAPZ).

Temporal distribution: May, June, July, October, and December.

Hosts: *Oryza sativa* L. (rice), *Asparagus officinalis* L. (asparagus) (EAPZ), and *Jatropha curcas* L. (Barbados nut) (Alonso and Lezcano 2014).

Recorded in Fabaceae (Maes and Goellner-Scheiding 1993).

***Chelinidea tabulata* (Burmeister, 1835)**

**Distribution:** Comayagua and Francisco Morazán.

**Notes:** Specimens examined: 6 (CEEF).

Temporal distribution: May and October.

Hosts: *Opuntia pilifera* F.A.C. Weber and *Opuntia imbricata* (Haw.) DC. (Brailovsky et al. 1994).

***Cimolus vitticeps* Stål, 1862**

**Distribution:** Distribution in Honduras unknown (Packauskas 2010).

**Notes:** Temporal distribution: Unknown.

Hosts: Unknown.

***Diactor bilineatus* (F., 1803)**

**Distribution:** Atlántida.

**Notes: NEW COUNTRY RECORD**

Number of specimens: 2 (CURLA).

Temporal distribution: July.

Hosts: *Passiflora edulis* Sims (passion fruit) (Oliveira and Frizzas 2014).

Known distribution: Brazil (Packauskas 2010).



It is considered one of the most important pests of passion fruit in Brazil (Oliveira and Frizzas 2014).

### ***Holhymenia histrio* (F., 1803)**

**Distribution:** Atlántida, Francisco Morazán, and Olancho.

**Notes:** NEW COUNTRY RECORD

Specimens examined: 9 (EAPZ)

Temporal distribution: February–July.

Hosts: *Passiflora edulis* Sims (passion fruit) (EAPZ), *Passiflora nitida* Kunth, *Passiflora coccinea* Aublet, *Passiflora gibertii* Brown, and *Passiflora alata* Curtis (Baldin and Boiça 1999).

Known distribution: Argentina, Brazil, Colombia, French Guiana, Nicaragua, Paraguay, Suriname, and Uruguay (Packauskas 2010).

This species is considered an important pest of passion fruit in Brazil (Baldin and Boiça 1999).

### ***Hypselonotus fulvus* (De Geer, 1773)**

**Distribution:** Atlántida and La Paz.

**Notes:** Specimens examined: 5 (CURLA).

Temporal distribution: August–September.

Hosts: *Zea mays* L. (corn), *Phaseolus vulgaris* L. (beans) (Passoa 1983), (Maes and Goellner-Scheiding 1993); *Ananas comosus* L. (pineapple) (Arellano et al. 2015), *Psidium guajava* L., *Persea americana* Miller (avocado), and *Gossypium hirsutum* L. (cotton) (Pires et al. 2013).

### ***Hypselonotus interruptus* Hahn, 1833**

**Distribution:** Atlántida, Choluteca, Comayagua, Cortés, El Paraíso, Francisco Morazán, Gracias a Dios, and Yoro.

**Notes:** Specimens examined: 102 (CEEF, CURLA, EAPZ, UNAH).

Temporal distribution: February–July.

Hosts: *Eupatorium* sp., *Rubus adenotrichos* Schltld., *Gossypium hirsutum* L. (cotton),

Hosts: *Litchi chinensis* Sonn., *Oryza sativa* L. (Rice) (Passoa 1983); *Ananas comosus* L. (pineapple) (Arellano et al. 2015); *Citrus limon* (L.) Osbeck (lemon) (USDA 2015); *Casearia sylvestris* Swartz, *Campomanesia xanthocarpa* (Mart.) O. Berg., *Myrciaria rivularis* (Cambess) O. Berg, *Acacia meanrsii* De Wild., *Caesalpinia pluviosa* DC., and *Cupressus macrocarpa* Hart. (cypress) (Thum and Costa 1997).

### ***Hypselonotus lineatus* Stål, 1862**

**Distribution:** Atlántida and Francisco Morazán.

**Notes:** Specimens examined: 9 (CURLA, EAPZ).

Temporal distribution: January, May–July.

Hosts: *Annona reticulata* L. (custard–apple), *Glycine max* (L.) Merr. (soy), *Sorghum halepense* (L.) Pers. (Johnson grass) (EAPZ), and *Jatropha curcas* L. (Barbados nut) (Alonso and Lezcano 2014).

### ***Hypselonotus punctiventris* Stål, 1862**

**Distribution:** Atlántida, Choluteca, Comayagua, Copán, Francisco Morazán, and Yoro.

**Notes:** Specimens examined: 119 (CEEF, EAPZ).

Temporal distribution: April–July and September–October.

Hosts: *Sesamum indicum* L. (sesame), *Sorghum halepense* (L.) Pers. (Johnson grass) (EAPZ); *Ipomoea batatas* (L.) Lam. (sweet potato), *Zea mays* L. (corn), *Coffea arabica* L. (coffee), *Citrus* sp. (Passoa 1983), and *Cirsium vulgare* (Savi) Tenore (Chordas et al. 2011).

### ***Leptoglossus brevirostris* Barber, 1862**

**Distribution:** Olancho.

**Notes:** **NEW COUNTRY RECORD**

Specimens examined: 1 (EAPZ).

Temporal distribution: August.

Hosts: *Zea mays* L. (corn) (EAPZ) and *Phoradendron leucarpum* (Raf.) Reveal & M. C. Johnst. (Whittaker 1984)

Known distribution: Costa Rica, Mexico, and United States (Packauskas 2010).



***Leptoglossus cinctus* (Herrich-Schäffer, 1836)**

**Distribution:** Atlántida, Comayagua, Francisco Morazán, and Yoro.

**Notes:** Specimens examined: 39 (CEEF, CURLA, EAPZ, UNAH).

Temporal distribution: October–November.

Hosts: *Citrus* sp., *Psidium guajava* L. (guava) (EAPZ), *Cereus* sp., and *Opuntia* sp. (Maes and Goellner-Scheiding 1993).

***Leptoglossus concolor* (Walker, 1871)**

**Distribution:** Atlántida, Comayagua, Francisco Morazán, Gracias a Dios, and Yoro.

**Notes:** Specimens examined: 17 (EAPZ).

Temporal distribution: May, July, September, and December.

Hosts: *Anacardium occidentale* L. (cashew), *Bixa orellana* L., *Psidium guajava* L., and *Litchi chinensis* Sonn. (Mitchell 2000).

***Leptoglossus crassicornis* (Dallas, 1852)**

**Distribution:** Yoro.

**Notes: NEW COUNTRY RECORD**

Specimens examined: 3 (CURLA).

Temporal distribution: August.

Hosts: *Harrisia pomanensis* (F.A.C. Weber ex K. Schum.) Britton & Rose, *Opuntia anacantha* Speg., *Opuntia elata* Link & Otto ex Salm-Dyck, *Opuntia discolor* Britton & Rose, *Opuntia sulphurea* Gillies ex Salm-Dyck, *Opuntia paraguayensis* K. Schum., *Opuntia quimilo* K. Schum., and *Opuntia ficus-indica* (L.) Mill. (Coscarón and Pall 2015).

Known distribution: Argentina, Bolivia, Colombia, Paraguay, and Uruguay (Packauskas 2010). This is the first record for Central America of a species previously believed to be restricted to South America. Further research is needed to clarify the status of this species.

***Leptoglossus gonagra* (F., 1775)**

**Distribution:** Choluteca, Comayagua, Francisco Morazán, and Olancho.

**Notes:** Specimens examined: 40 (CEEF, EAPZ)

Temporal distribution: January, May–July, September, and October.

Hosts: *Luffa cylindrica* (L.) M. Roem., *Passiflora edulis* Sims (Passion fruit), *Citrullus lanatus* (Thunb.) Matsum. & Nakai (watermelon) (EAPZ); *Momordica charantia* L. (bitter melon), *Citrus sinensis* (L.) Osbeck (oranges), *Punica granatum* L. (grenade), *Mangifera indica* L. (mango), *Nicotiana tabacum* L. (tobacco), *Cucumis melo* L. (sweet melon), *Citrus paradisi* Macfadyen (grapefruit) (Mitchell 2000), and *Jatropha curcas* L. (Barbados nut) (Alonso and Lezcano 2014).

### ***Leptoglossus lineosus* (Stål, 1862)**

**Distribution:** Atlántida and Francisco Morazán.

**Notes:** NEW COUNTRY RECORD

Specimens examined: 2 (EAPZ, UNAH).

Temporal distribution: March–April.

Hosts: *Cucurbita* sp. (Mitchell 2000).

Known distribution: Mexico (Packauskas 2010).

### ***Leptoglossus oppositus* (Say, 1832)**

**Distribution:** Atlántida and Francisco Morazán.

**Notes:** Specimens examined: 1 (EAPZ).

Temporal distribution: May–July.

Hosts: *Pinus* sp., *Cucumis sativus* L. (cucumber) (Mitchell 2000); *Helianthus* sp., *Cucurbita* sp., *Citrullus lanatus* (Thunb.) Matsum. & Nakai (watermelon), *Carya* sp., *Yucca* sp., *Gossypium* sp., *Morus* sp., *Psidium guajava* L., *Zea mays* L. (corn), *Prunus* sp., *Pyrus* sp., *Coffea arabica* L. (coffee), *Datura* sp., and *Solanum lycopersicum* L. (tomato) (Maes and Goellner-Scheiding 1993).

Maes and Goellner-Scheiding (1993) recorded *Bicyrtes quadrifasciatus* Lepeletier (Hymenoptera: Crabronidae) as a natural enemy in Nicaragua. Adults are known to be parasitized by *Trichopoda pennipes* (F.) (Diptera: Tachinidae) (Mitchell 2000).

### ***Leptoglossus zonatus* (Dallas, 1852)**

**Distribution:** Atlántida, Choluteca, Comayagua, Copán, Cortés, El Paraíso, Francisco Morazán, Gracias a Dios, Islas de la Bahía, Lempira, Olancho, and Yoro.

**Notes:** Specimens examined: 195 (CEEF, CURLA, EAPZ, UNAH).



Temporal distribution: Year long.

Hosts: *Luffa cylindrica* Miller, *Passiflora edulis* Sims (passion fruit), *Asparagus officinalis* L. (asparagus), *Citrus sinensis* (L.) Osbeck (orange), *Solanum lycopersicum* L. (tomato), *Zea mays* L. (corn), *Punica granatum* L. (grenada), *Anacardium occidentale* L. (cashew), *Psidium guajava* L., *Oryza sativa* L. (rice), *Solanum tuberosum* L. (potato) (EAPZ); *Cucurbita* sp., *Triadica sebifera* (L.) Small (Chinese tallow), *Sorghum* sp. (sorghum), *Schizocarpum reflexum* Rose, *Chilopsis linearis* (Cav.) Sweet, *Jatropha curcas* L. (Barbados nut), *Actinocheita filicina* (DC.) F. A. Barkley (Mitchell 2000); *Helianthus* sp., *Crescentia* sp., *Hylocereus* sp., *Schizocarpum* sp., *Cucumis* sp., *Phaseolus* sp., *Persea* sp., *Gossypium* sp., *Azadirachta indica* A. Juss., *Musa* sp., *Sesamum indicum* L. (sesame), and *Coffea arabica* L. (coffee) (Maes and Goellner-Scheiding 1993).

*Leptoglossus zonatus* is one of the most important and abundant pest species of coreids in Honduras. This species is parasitized by wasps of the genera *Geyon*, *Ooencyrtus*, *Anastatus* and *Neorileya*. *Beauveria bassiana* and *Metarhizium anisopliae* have been used successfully as a control (Mitchell 2000).

### ***Leptoscelis quadrisignatus* (Distant, 1881)**

**Distribution:** Distribution in Honduras unknown (Packauskas 2010).

**Notes:** Temporal distribution: Unknown.

Hosts: Unknown.

### ***Lycambes andicola* (Breddin, 1903)**

**Distribution:** Atlántida, Comayagua, and Francisco Morazán.

**Notes: NEW COUNTRY RECORD**

Specimens examined: 4 (CEEf, CURLA, EAPZ, UNAH).

Temporal distribution: July–September.

Known distribution: Bolivia (Packauskas 2010).

### ***Machtima mexicana* Stål, 1870**

**Distribution:** Francisco Morazán.

**Notes: NEW COUNTRY RECORD**

Specimens examined: 7 (EAPZ, UNAH).

Temporal distribution: May, June, and August.

Known distribution: Mexico and Panama (Packauskas 2010).

### ***Madura perfida* Stål, 1862**

**Distribution:** Distribution in Honduras unknown (Packauskas 2010).

**Notes:** Temporal distribution: Unknown.

Hosts: *Zea mays* L. (corn) (Gibson and Carrillo 1959).

### ***Melucha phyllocnemis* (Burmeister, 1835)**

**Distribution:** Francisco Morazán.

**Notes:** NEW COUNTRY RECORD

Specimens examined: 1 (UNAH).

Temporal distribution: July.

Hosts: *Neurolaena lobata* (L.) R.Br. ex Cass., *Mikania scandens* (L.) Willd., and *Verbesina* sp. (Barcellos et al. 2008).

Known distribution: Bolivia, Brazil, Mexico, Colombia, and Paraguay (Packauskas 2010).

### ***Melucha quadrivittis* Stål, 1862**

**Distribution:** Copán, Francisco Morazán, and Olancho.

**Notes:** Specimens examined: 3 (UNAH).

Temporal distribution: May, September, and November.

### ***Mozena lineolata* (Herrich-Schäffer, 1842)**

**Distribution:** Atlántida, Francisco Morazán, and Olancho (Brailovsky and Barrera 2014).

**Notes:** Specimens examined: 128 (CEEF, CURLA, EAPZ, UNAH).

Temporal distribution: May–October.

Hosts: *Prosopis* sp. and *Atriplex* sp.



***Mozena lunata* (Burmeister, 1835)**

**Distribution:** Atlántida, Comayagua, Francisco Morazán, and Yoro.

**Notes:** NEW COUNTRY RECORD

Specimens examined: 152 (CEEF, CURLA, EAPZ, UNAH).

Temporal distribution: May–November.

Hosts: *Malus domestica* Borkh., *Cassia* sp. (EAPZ); *Acacia farnesiana* L. Willd. (Ward et al. 1977), *Acacia amentacea* DC., and *Prosopis glandulosa* Torr. (Brailovsky et al. 1995).

Known distribution: Costa Rica, Guatemala, Mexico, Nicaragua, and United States (Packauskas 2010).

***Mozena lurida* (Dallas, 1852)**

**Distribution:** Distribution in Honduras unknown (Packauskas 2010).

**Notes:** Temporal distribution: Unknown.

Hosts: Unknown.

***Mozena lutea* (Herrich-Schaeffer, 1840)**

**Distribution:** Distribution in Honduras unknown (Packauskas 2010).

**Notes:** Temporal distribution: Unknown.

Hosts: Unknown.

***Nematopus lepidus* Stål, 1862**

**Distribution:** Atlántida, Olancho, and Yoro.

**Notes:** NEW COUNTRY RECORD

Specimens examined: 11 (EAPZ).

Temporal distribution: March–August.

Hosts: *Persea americana* Miller (avocado) (EAPZ).

Known distribution: Mexico and Panama (Packauskas 2010).

***Pachylis nervosus* Dallas, 1852**

**Distribution:** Atlántida, Comayagua, El Paraíso, Francisco Morazán, Islas de la Bahía, Olancho, and Yoro.

**Notes:** Specimens examined: 255 (CEEF, CURLA, EAPZ, UNAH).

Temporal distribution: Year long.

Hosts: *Mimosa* sp. and *Prosopis laevigata* (Willd.) M. C. Johnst. (Brailovsky 1995).

***Paryphes anceps* Horvath, 1913**

**Distribution:** Cortés (Horvath 1913)

**Notes:** Temporal distribution: Unknown.

Hosts: Unknown.

***Paryphes flavocinctus* Stål, 1860**

**Distribution:** Atlántida, Francisco Morazán, and Yoro.

**Notes:** Specimens examined: 8 (CURLA, EAPZ)

Temporal distribution: April–July.

Hosts: *Citrullus lanatus* (Thunb.) Matsum. & Nakai (watermelon) and *Gurania* sp. (Gilbert 1991).

***Phthia lunata* (F., 1787)**

**Distribution:** Atlántida, Comayagua, and Cortés.

**Notes: NEW COUNTRY RECORD**

Specimens examined: 5 (EAPZ).

Temporal distribution: June, August, and September.

Hosts: *Capsicum annuum* L. (sweet pepper) (EAPZ); *Citrullus lanatus* (Thunb.) Matsum. & Nakai (watermelon), and *Cucurbita* sp. (Maes and Goellner-Scheiding 1993).

Known distribution: Argentina, Brazil, Colombia, Costa Rica, Cuba, Mexico, Panama, Puerto Rico, and Suriname (Packauskas 2010).

This species is reported as a vector of *Phytophthora* sp. (Godoi et al. 2002).



***Peranthus longicornis* (Dallas, 1852)**

**Distribution:** Atlántida.

**Notes:** NEW COUNTRY RECORD

Specimens examined: 1 (EAPZ).

Temporal distribution: April.

Known distribution: Brazil (Packauskas 2010).

***Phthiacnemia picta* (Drury, 1773)**

**Distribution:** Atlántida, Comayagua, Choluteca, El Paraíso, and Francisco Morazán.

**Notes:** Specimens examined: 38 (CEEF, CURLA, EAPZ, UNAH).

Temporal distribution: May–July, August, and September.

Hosts: *Ipomoea batatas* (L.) Lam. (sweet potato), *Solanum tuberosum* L. (potato), *Solanum lycopersicum* L. (tomato) (EAPZ); *Pisum* sp., *Phaseolus vulgaris* L. (beans), *Trifolium* sp., *Vicia* sp., *Vigna* sp., *Stellaria* sp., *Citrullus lanatus* (Thunb.) Matsum. & Nakai (watermelon), *Cucurbita pepo* L. (pumpkin), *Cucumis sativus* L. (cucumber), *Momordica* sp., *Passiflora edulis* Sims (passion fruit), *Punica granatum* L. (grenade), *Sorghum* sp., *Zea mays* L. (corn), *Oryza sativa* L. (rice), *Coffea arabica* L. (coffee), *Sesamum indicum* L. (sesame), *Helianthus annuus* L. (sunflower), and *Nicotiana tabacum* L. (tobacco) (Maes and Goellner-Scheiding 1993).

This species is considered an important pests of tomato in Brazil (Da Silva et al. 2003).

***Piezogaster auriculatus* (Stål, 1862)**

**Distribution:** Atlántida, Comayagua, Francisco Morazán, and Yoro.

**Notes:** Specimens examined: 54 (CEEF, CURLA, EAPZ, UNAH).

Temporal distribution: May–October.

Hosts: *Citrus sinensis* (L.) Osbeck (oranges) and *Pachyrhizus* sp. (Dealy 2000).

***Plapigus circumcinctus* Stål, 1860**

**Distribution:** Atlántida, Comayagua, Francisco Morazán, and Yoro.

**Notes:** Specimens examined: 66 (CEEF, CURLA, EAPZ, UNAH).

Temporal distribution: May–July and October–December.

Hosts: *Coffea arabica* L. (coffee) (Maes and Goellner-Scheiding 1993).

***Possaniella oblata* Brailovsky, 1999**

**Distribution:** Atlántida.

**Notes:** NEW COUNTRY RECORD

Specimens examined: 1 (CURLA).

Temporal distribution: May.

Known distribution: Brazil (Packauskas 2010).

***Romoniella perfecta* Brailovsky & Barrera, 2001**

**Distribution:** Atlántida, Comayagua, Olancho, and Yoro.

**Notes:** Specimens examined: 9 (CEEF, EAPZ, CURLA).

Temporal distribution: February, October, and November.

***Sagotylus confluens* (Say, 1832)**

**Distribution:** Atlántida, Comayagua, Choluteca, Cortés, El Paraíso, Francisco Morazán, Islas de la Bahía, Olancho, and Yoro.

**Notes:** Specimens examined: 196 (CEEF, CURLA, EAPZ, UNAH).

Temporal distribution: Year long.

Hosts: *Oryza sativa* L. (rice), *Zea mays* L. (corn) (Passoa 1983), *Canavalia ensiformis* (L.) DC. (EAPZ); *Ricinus communis* L. (Valdés-Rodríguez et al. 2015), and *Senna obtusifolia* L. H .S. Irwin & Barneby (Palmer and Pullen 2000).

This species is considered one of the main pests of fig trees in Mexico feeding on terminal shoots and new leaves. Due to its large size, the latex loss is high and its excreta favor the establishment of fungi on the terminal buds causing them to dry (Valdés-Rodríguez et al. 2015).

***Salamancaniella alternata* (Dallas, 1852)**

**Distribution:** Atlántida and Comayagua.

**Notes:** Specimens examined: 7 (CURLA, EAPZ).

Temporal distribution: May, July, August–November.



Hosts: *Jatropha curcas* L. (Barbados nut) (Alonso and Lezcano 2014).

***Savius jurgiosus* (Stål, 1862)**

**Distribution:** Francisco Morazán.

**Notes:** NEW COUNTRY RECORD

Specimens examined: 2 (EAPZ).

Temporal distribution: October.

Hosts: *Baltimora* sp., *Buddleja sessiliflora* Kunth, *Prosopis* sp., and *Ziziphus* sp.

Known distribution: Costa Rica, Guatemala, Mexico, and United States (Packauskas 2010).

***Sephina limbata* Stål, 1862)**

**Distribution:** Atlántida and Comayagua.

**Notes:** Specimens examined: 13 (CEEF, CURLA)

Temporal distribution: January–April.

Hosts: Unknown.

***Serranoniella amblysa* Brailovsky & Barrera, 2001**

**Distribution:** Comayagua and Yoro.

**Notes:** NEW COUNTRY RECORD

Specimens examined: 22 (CEEF).

Temporal distribution: June, September–December.

Known distribution: Brazil (Packauskas 2010).

***Spartocera fusca* (Thunberg, 1783)**

**Distribution:** Comayagua and Yoro.

**Notes:** Specimens examined: 71 (CEEF, CURLA, EAPZ, UNAH).

Temporal distribution: Year long.

Hosts: *Ipomoea batatas* (L.) Lam. (sweet potato), *Gossypium* sp. (cotton), *Solanum lycopersicum* L. (tomato), *Solanum americanum* Miller, *Physalis peruviana* L. (Maes

and Goellner-Scheiding 1993); *Solanum tuberosum* L. (potato) (Passoa 1983), *Solanum nigrum* L. (Mitchell 2000), and *Capsicum annum* L. (Mitchell 2000).

Notes: *Sarcophaga sternodontis* Towns (Diptera: Tachinidae) is a known parasite of *S. fusca* in Nicaragua (Maes and Goellner-Scheiding 1993).

### ***Staluptus marginalis* (Burmeister, 1835)**

**Distribution:** Comayagua, Francisco Morazán, and Yoro.

**Notes: NEW COUNTRY RECORD**

Specimens examined: 39 (CEEF, EAPZ).

Temporal distribution: May, July, September, and October.

Hosts: *Sorghum halepense* (L.) Pers. (Johnson grass) (EAPZ); and *Cajanus cajan* (L.) Millsp. (Maes and Goellner-Scheiding 1993).

Known distribution: Guatemala and Mexico (Packauskas 2010).

### ***Thasopsis formidabilis* (Distant, 1893)**

**Distribution:** Atlántida, Comayagua, Francisco Morazán, and Santa Bárbara

**Notes: NEW COUNTRY RECORD**

Specimens examined: 13 (CURLA, EAPZ).

Temporal distribution: February, May, and September.

Known distribution: Costa Rica and Panama (Packauskas 2010).

### ***Thasus acutangulus* (Stål, 1859)**

**Distribution:** Atlántida and Yoro.

**Notes:** Specimens examined: 2 (CURLA, EAPZ).

Temporal distribution: December.

Hosts: *Prosopis velutina* Wooton (Torre-Bueno 1945), and *Pithecellobium* sp. (EAPZ).

### ***Zicca rubricator* (F., 1803)**

**Distribution:** Unknown (Brailovsky and Cadena 1992).

**Notes:** Temporal distribution: Unknown.



Hosts: *Schaueria calycobractea* Hilsenbeck & D. L. Marshall, *Chamissoa altissima* (Jacq.) Kunth and *Phytolacca rivinoides* Kunth & C. D. Bouché (Brailovsky and Cadena 1992).

**Zicca taeniola (Dallas, 1852)**

**Distribution:** Atlántida, El Paraíso, Francisco Morazán, and Olancho.

**Notes:** Specimens examined: 32 (CURLA, EAPZ).

Temporal distribution: December.

Hosts: *Glycine max* (L.) Merr. (soy), *Solanum tuberosum* L. (potato), *Zea mays* L. (corn), *Sechium edule* (Jacq.) Sw., *Daucus carota* L. (carrot) (EAPZ), and *Capsicum annum* L. (Maes and Goellner-Scheiding 1993).

**Discussion**

A total of 2,036 specimens were examined. 68 species are now known from Honduras with 19 being new country records (Table 1). Seventeen species were only found recorded in literature.

Table 1. New country records of Coreidae for Honduras		
Species	Specimens examined	Collection
<i>Anasa trilineata</i> Stål	11	EAPZ
<i>Camptischium clavipes</i> (F.)	8	EAPZ
<i>Diactor bilineatus</i> (F.)	2	CURLA
<i>Holhymenia histrio</i> (F.)	9	EAPZ
<i>Leptoglossus brevirostris</i> Barber	1	EAPZ
<i>Leptoglossus lineosus</i> (Stål)	2	EAPZ, UNAH
<i>Leptoglossus crassicornis</i> (Dallas)	3	CURLA
<i>Lycambes andicola</i> Breddin	4	All collections
<i>Machtima mexicana</i> Stål	7	EAPZ y UNAH
<i>Melucha phyllocnemis</i> (Burmeister)	1	UNAH
<i>Mozena lunata</i> (Burmeister)	152	All collections
<i>Nematopus lepidus</i> Stål	11	EAPZ
<i>Peranthus longicornis</i> (Dallas)	1	EAPZ
<i>Phthia lunata</i> (F.)	5	EAPZ
<i>Possaniella oblata</i> Brailovsky	1	CURLA

<i>Savius jurgiosus</i> (Stål)	2	EAPZ
<i>Serranoniella amblysa</i> Brailovsky & Barrera	22	CEEF
<i>Staluptus marginalis</i> (Burmeister)	39	EAPZ, CEEF
<i>Thasopsis formidabilis</i> (Distant)	13	EAPZ, CURLA

Many of the species found in Honduras are known only from one department. We suspect this is not a real reflection of the species' distribution but collecting bias. For nine of the species there is no information on where in Honduras they occur.

## Acknowledgements

We thank the curators at each of the institutions: Karla Cantarero (UNAH), Carlos Salgado (CURLA), and Oscar Leverón (CURLA). This research was funded by the Department of Agricultural Science and Production at Zamorano University.

## References

- Alonso O, Lezcano JC (2014) Artrópodos asociados a *Jatropha curcas* Linnaeus. Funciones y estrategia para su manejo. Past. Forr 37 (1): 3-16.
- Alston DG, Barnhill JV (2008) Squash Bug (*Anasa tristis*). Utah Pests Fact Sheet ENT 120 (08): 1-6.
- Arellano G, Vergara C, Bello S (2015) Plagas entomológicas y otros artrópodos en el cultivo de la piña (*Ananas comosus* var. *comosus* (L.) Merr., Coppens & Leal) en Chanchamayo y Satipo, departamento de Junín, Perú. Ecol. Apl 14 (2): 175-189. <https://doi.org/10.21704/rea.v14i1-2.94>
- Báez J, Cervantes L (2014) Estados inmaduros y fenología de *Catorhintha apicalis scrutator* (Hemiptera: Heteroptera: Coreidae) en Michoacán, México. Rev. Biol. Trop 62 (2): 579-587. <https://doi.org/10.15517/rbt.v62i2.11783>
- Baldin E, Boiça A (1999) Desenvolvimento de *Holhymenia histrio* (Fabr.) (Hemiptera: Coreidae) em frutos de cinco genótipos de maracujazeiro (*Passiflora* spp.). An. Soc. Entomol Bras 28 (3): 421-427. <https://doi.org/10.1590/S0301-80591999000300006>
- Barcellos A, Schmidt L, Brailovsky H (2008) Abundance and Species Richness of Coreoidea (Hemiptera: Heteroptera) from Parque Estadual do Turvo. Southern Brazil. Neotrop. Entomol 37 (4): 406-412. <https://doi.org/10.1590/S1519-566X2008000400008>
- Brailovsky H (1985) Revisión del género *Anasa* Amyot-Serville (Hemiptera, Heteroptera, Coreidae, Coreinae, Coreini). Monog. Inst. Biol. Univ. Auton. Mex 2: 1-226.
- Brailovsky H (1995) Revisión del complejo "Cebrenis" (Hemiptera-Heteroptera-Coreidae-Coreinae-Coreini). Publ. Esp. Inst. Biol. Univ. Auton. Mex 15: 1-124.
- Brailovsky H (2013) Description of one new species of *Cebrenistella*, and key to the known species (Hemiptera: Heteroptera: Coreidae). Dtsch. Entomol. Z 60: 209-212.
- Brailovsky H (2016) A new species of *Anasa* from Honduras (Hemiptera: Heteroptera: Coreidae). Entomologica Americana 122 (1-2): 31-36. <https://doi.org/10.1664/15-RA-032>



- Brailovsky H, Barrera E (2014) Revisional notes on the genus *Melucha* (Hemiptera, Heteroptera, Coreidae). Dtsch. Entomol. Z 61: 15-22. <https://doi.org/10.3897/dez.61.7048>
- Brailovsky H, Cadena A (1992) Revisión del género *Zicca* (Hemiptera- Heteroptera- Coreidae-Coreinae-Coreini). Publ. Esp. Inst. Biol. Univ. Auton. Mex 9: 9-101.
- Brailovsky H, Garcia M (1987) Revisión del género *Catorhintha* Stål (Hemiptera- Heteroptera-Coreidae-Coreinae-Coreini). Monog. Inst. Biol. Univ. Auton. Mex 4: 1-148.
- Brailovsky H, Barrera E, Mayorga C, Ortega G (1994) Estadios ninfales de los Coreidos del Valle de Tehuacán, Puebla (Hemiptera-Heteroptera) *Chelinidea staffilesi*, *C. tabulata* y *Narnia femorata* . An. Inst. Biol. Univ. Auton. Mex 65 (2): 250-258.
- Brailovsky H, Mayorga C, Ortega G, Barrera E (1995) Estadios ninfales de los Coreidos del Valle de Tehuacán, Puebla, México (Hemiptera-Heteroptera) II. Especies asociadas a Huizacheras (*Acacia* spp.) y Mezquiteras (*Prosopis* spp.): *Mozena lunata*, *Pachylis hector*, *Savius jurgiosus* y *Thasus gigas* . An. Inst. Biol. Univ. Auton. Mex 66 (1): 57-80.
- Cervantes L, Báez J, Brailovsky H (2014) Estados inmaduros de coreidos (Hemiptera: Heteroptera: Coreidae: Coreinae: Coreini: Chelinideini) de Baja California, México. Rev. Mex. Biod 85: 741-752. <https://doi.org/10.7550/rmb.43261>
- Chordas SW, Tumilson R, Robison HW, Kremers J (2011) Twenty Three True Bug State Records for Arkansas, with Two for Ohio, U.S.A. J. Ark. Acad. Sc 65 (1): 153-159.
- Coscarón M, Pall J (2015) The Tribe Anisoscelini (Hemiptera: Heteroptera, Coreidae) in Argentina. Zootaxa 4033 (3): 411-426. <https://doi.org/10.11646/zootaxa.4033.3.6>
- Da Silva RA, Carvalho GS, Flores PS (2003) Morfologia externa dos adultos de *Phthia picta* (Drury, 1770) (Hemiptera, Coreidae). Bol. Veg. Plagas 29: 249-253.
- Dealy B (2000) A Revision of the genus *Piezogaster* Amyot & Serville, (Heteroptera: Coreidae: Nematopodini) and the description of two new species. Fort Hays State University, Hays
- Evans GA, Halbert SE (2007) A Checklist of the Aphids of Honduras (Hemiptera: Aphididae). Fla. Entomol 80 (3): 518-523. [https://doi.org/10.1653/0015-4040\(2007\)90\[518:ACOTAO\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2007)90[518:ACOTAO]2.0.CO;2)
- Gibson WW, Carrillo JL (1959) Lista de insectos en la colección entomológica de la oficina de estudios especiales, SAG. Inst. Nac. Inv. Agr. Fol. Misc 9: 1-254.
- Gilbert LE (1991) *Anguria* y *Gurania* (Cucurbitaceae) (Pepinos de Monte). In: Janzen DH (Ed.) Historia natural de Costa Rica. Editorial de la Universidad de Costa Rica, San José.
- Godoi MM, Serrano MG, Teixeira MM, Camargo EP (2002) A PCR–Based Survey on *Phytomonas* (Euglenozoa: Trypanosomatidae) in Phytophagous Hemipterans of the Amazon Region. J. Eukaryot. Microbiol 49 (4): 275-279. <https://doi.org/10.1111/j.1550-7408.2002.tb00370.x>
- Grimm C, Maes JM (1997) Insectos asociados al cultivo de tempate (*Jatropha crucas*) en el Pacífico de Nicaragua. Rev. Nica. Ent 42: 16-34.
- Henry TJ (2009) Biodiversity of Heteroptera . In: Footitt E, Adler P (Eds) Insect Biodiversity: Science and Society. Wiley- Blackwell, Oxford.
- Hernández LM, Pinzón JM (2015) Primer Reporte de *Piezogaster odiosus* Stål y *Acanthocephala alata* Burmeister (Hemiptera: Coreidae) como plagas de *Annona muricata* L. Southwest. Entomol 40 (3): 665-668. <https://doi.org/10.3958/059.040.0323>
- Horvath G (1913) Revisio critica generis *Paryphes* Burm. et affinium. Annales Musei Nationalis Hungarici 11: 344-373.



- King AB, Saunders JS (1984) The Invertebrate Pests of Annual Food Crops in Central America: A Guide to Their Recognition and Control. Overseas Development Administration, London.
- Lerma L, Rojas IE, Velasco NR, Figueroa F (1986) Guía práctica para la identificación y manejo de las principales plagas en cultivos frutales. Acta Agron 36 (3): 68-74.
- Lucas K, Maggi J, Yagual M (2010) Creación de una empresa de producción, comercialización y exportación de tomate de árbol en el área de Sangolquí, provincia de Pichincha. BSc dissertation. Escuela Politécnica del Litoral, Guayaquil.
- Maes JM, Goellner-Scheiding U (1993) Catálogo de los Coreidae (Heteroptera) de Nicaragua. Rev. Nica. Ent 25: 1-19.
- McPherson JE, Packauskas RJ, Sites RW, Taylor SJ, Bundy CS, Bradshaw JD, Mitchell PL (2011) Review of *Acanthocephala* (Hemiptera: Heteroptera: Coreidae) of America north of Mexico with a key to species. Zootaxa 2835: 30-40.
- Mitchell PL (2000) Leaf-footed bugs (Coreidae). In: Schaefer CW, Panizzi AR (Eds) Heteroptera of Economic Importance. CRC Press, Boca Raton, FL.
- Oliveira CM, Frizzas MR (2014) Principais Pragas do Maracujazeiro Amarelo (*Passiflora edulis* f. *flavicarpa* Degener) e seu Manejo. EMBRAPA Cerr. Doc 323: 1-44.
- Packauskas R (2010) Catalog of the Coreidae, Or Leaf-footed Bugs, of the New World. Fort Hays Studies
- Palmer WA, Pullen KR (2000) The Phytophagous Arthropods Associated with *Senna obtusifolia* (Caesalpiniaceae) in Mexico and Honduras and Their Prospects for Utilization for Biological Control. Biol. Con 20 (1): 76-83. <https://doi.org/10.1006/bcon.2000.0879>
- Passoa S (1983) Lista de los insectos asociados con los granos básicos y otros cultivos selectos en Honduras. Ceiba 25 (1): 7-18.
- Pires EM, Nogueira RM, da Silva CJ, Pelissari F, Ferreira JAM, Soares MA (2013) New Sucking Coreids Species in *Psidium guajava* . Sci. Elec. Arch 4: 31-35.
- Thum AB, Costa EC (1997) Coreidae (Heteroptera) associados a espécies florestais. Cienc. Flo 7 (1): 27-31.
- Torre-Bueno JR (1945) Random notes on *Thasus acutangulus* . Bull. Brooklyn Ent. Soc 40: 83.
- Turnbow RH, Cave RD, Kingsolver JM (2003a) An Annotated Checklist of the Bruchidae of Honduras. Ceiba 44 (2): 269-278.
- Turnbow RH, Cave RD, Thomas MC (2003b) A list of the Cerambycidae of Honduras, with additions of previously unrecorded species. Ceiba 44 (1): 1-43.
- USDA (2015) Risk Assessment for the Importation of Fresh Lemon (*Citrus limon* (L.) Burm. f.) Fruit from Northwest Argentina into the Continental United States. USDA, Raleigh, N.C..
- Valdés-Rodríguez OA, Pérez-Vázquez A, Palacios-Wassenaar OM (2015) Insectos plaga en cultivo asociado de *Ricinus communis* y *Moringa oleifera* en el centro de Veracruz, México. Rev. Mex. Cienc. Agr. Pub. Esp 11: 2233-2239.
- Ward CR, O'Brien CW, O'Brien LB, Foster DE, Huddleston EW (1977) Annotated Checklist of New World Insects Associated with *Prosopis* (Mesquite). U.S. Dep. Agric. Tech. Bull 1557: 1-115.
- Whittaker PL (1984) The insect fauna of mistletoe (*Phoradendron tomentosum*, Loranthaceae) in southern Texas. Southwest. Nat 29: 435-444. <https://doi.org/10.2307/3670996>